

M. V. JOHNS JR.

EFFICIENT ESTIMATION OF REGRESSION COEFFICIENTS  
IN TIME SERIES

BY

T. W. ANDERSON

TECHNICAL REPORT NO. 2

OCTOBER 1, 1970

PREPARED UNDER CONTRACT

N00014-67-A-0112-0030 (NR-042-034)

FOR THE OFFICE OF NAVAL RESEARCH

THEODORE W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS  
STANFORD, UNIVERSITY  
STANFORD, CALIFORNIA



EFFICIENT ESTIMATION OF REGRESSION COEFFICIENTS  
IN TIME SERIES

BY

T. W. ANDERSON  
Stanford University

TECHNICAL REPORT NO. 2

OCTOBER 1, 1970

PREPARED UNDER THE AUSPICES

OF

OFFICE OF NAVAL RESEARCH CONTRACT #N00014-67-A-0112-0030

THEODORE W. ANDERSON, PROJECT DIRECTOR

DEPARTMENT OF STATISTICS  
STANFORD UNIVERSITY  
STANFORD, CALIFORNIA

## 1. Introduction

This paper deals with estimating regression coefficients in the usual linear model. Let  $\tilde{y}$  be a T-component random vector with expected value

$$(1) \quad \tilde{E} \tilde{y} = \tilde{Z} \tilde{\beta} ,$$

where  $\tilde{Z}$  is a  $T \times p$  matrix of numbers and  $\tilde{\beta}$  is a p-component vector of parameters. (All vectors are column vectors.) For convenience we assume that the rank of  $\tilde{Z}$  is the number of columns,  $p$ . The covariance matrix of  $\tilde{y}$  is

$$(2) \quad \tilde{C}(\tilde{y}) = \tilde{C}(\tilde{y} - \tilde{Z} \tilde{\beta})(\tilde{y} - \tilde{Z} \tilde{\beta})' = \tilde{\Sigma} .$$

(Transposition of a vector or matrix is denoted by a prime.) Again for convenience we shall assume that  $\tilde{\Sigma}$  is positive definite. The problem is to estimate  $\tilde{\beta}$  on the basis of one observation on  $\tilde{y}$  when  $\tilde{Z}$  is known.

When  $\tilde{\Sigma}$  is known or is known to within a constant multiple, the Markov or Best Linear Unbiased Estimate (BLUE) is given by

$$(3) \quad \tilde{b} = (\tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z})^{-1} \tilde{Z}' \tilde{\Sigma}^{-1} \tilde{y} .$$

The least squares estimate is given by

$$(4) \quad \tilde{b}^* = (\tilde{Z}' \tilde{Z})^{-1} \tilde{Z}' \tilde{y} .$$

The covariance matrix of the Markov estimate is

$$(5) \quad \mathcal{L}(\tilde{b}) = (\tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z})^{-1} ;$$

the covariance matrix of the least squares estimate is

$$(6) \quad \mathcal{L}(\tilde{b}^*) = (\tilde{Z}' \tilde{Z})^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} (\tilde{Z}' \tilde{Z})^{-1} .$$

Both of the estimates are linear and unbiased.

The optimality property of the Markov estimate implies that  $\mathcal{L}(\tilde{b}^*) - \mathcal{L}(\tilde{b})$  is positive semidefinite; that is, any linear function of the Markov estimate has a variance no larger than the variance of that linear function of the least squares estimate. Since the least squares estimate can always be calculated, but the Markov estimate is unavailable if the covariance matrix  $\tilde{\Sigma}$  is now known to within a constant of proportionality, an interesting problem is to find under what conditions the least squares estimate is identical to the Markov estimate. It will be noted that they are identical when  $\tilde{\Sigma}$  is a multiple of the identity,  $I$ . The general answer is given by the following theorem:

Theorem 1. The least squares estimate (4) is identical to the best linear unbiased estimate (3) if and only if  $\tilde{Z} = \tilde{V}^* \tilde{C}$ , where the  $p$  columns of  $\tilde{V}^*$  are  $p$  linearly independent characteristic vectors of  $\tilde{\Sigma}$  and  $\tilde{C}$  is a nonsingular matrix.

The sufficiency of the condition was essentially given by myself in 1948 in the Skandinavisk Aktuarietidskrift [1]. In that paper I showed that if  $\tilde{y}$  is normally distributed, then the least squares

estimate is identical to the maximum likelihood estimate; under normality, of course, the maximum likelihood estimate is best linear unbiased. Watson [9], [10], G. S. Watson and Hannan [12] studied the efficiency of least squares estimates; the inequality given in the first two papers shows the necessity of the condition for  $p = 1$ . Magness and McGuire [6] rediscovered the condition, proving sufficiency and necessity. Watson [11] and Zyskind [13] have made more intensive studies and surveyed the literature.

A problem that is more explicitly and specially a time series problem occurs in the case where the residuals constitute a stationary stochastic process. The property  $\sigma_{st} = \sigma(s - t)$ , where  $\Sigma = (\sigma_{st})$  denotes stationarity in the wide sense. In general, the least squares estimate and the best linear unbiased estimate will be different. The characteristic vectors of  $\Sigma$  depend on the values of the serial or lag covariances and hence the best linear unbiased estimate depends on these parameters, which are generally unknown.

In this case we consider the covariance matrices of the estimates, normalize them suitably and identically, and consider the limits of them as  $T \rightarrow \infty$ . Grenander [4], Rosenblatt [7] in the Third Berkeley Symposium, and these two authors [5] found conditions for which the two limiting covariance matrices are the same. They did not indicate that their results were asymptotic analogues of the result for a finite sample, and the statement of their results and their methods of proof do not make it easy to see the relationship.

In this paper I shall prove the results for the finite-dimensional case and the limiting case in a similar fashion in order that the relationship between the results be clearer and that the asymptotic results be more easily understood. The emphasis here is on the linear algebra; the rigorous derivation of the limits, which is rather involved is omitted (but is given in Section 10.2 of [3]).

The method of proof is not the most direct for Theorem 1, because the proof uses covariance matrices instead of the structure of the estimates themselves. On the other hand, the asymptotic results must be derived in terms of the covariance matrices because the order of the observation vector increases, and thus the structure of the estimate changes. To obtain comparable proofs, covariance matrices must be used throughout. A by-product of my proof of the theorems is a different statement of the conditions of Grenander and Rosenblatt, which, I hope, is more enlightening than the original. Watson [9] related the two sets of results by considering the finite-sample case in the framework of the approach of Grenander and Rosenblatt.

## 2. The Finite-Sample Case.

We shall now proceed to prove Theorem 1 by considering the conditions for which the two covariance matrices, (5) and (6), are identical. To study this problem it will be convenient to transform the coordinate system in the  $T$ -dimensional space to the coordinate system defined by the characteristic vectors of the covariance matrix  $\Sigma$ . Let

$$(7) \quad \tilde{\Lambda} = \begin{pmatrix} \lambda_1 & 0 & \dots & 0 \\ 0 & \lambda_2 & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & \lambda_T \end{pmatrix},$$

where  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_T (> 0)$  are the characteristic roots of  $\tilde{\Sigma}$ . Let  $\tilde{V}$  be a  $T \times T$  matrix with columns as corresponding normalized characteristic vectors. These properties can be summarized in the two matrix equations

$$(8) \quad \tilde{\Sigma}\tilde{V} = \tilde{V}\tilde{\Lambda},$$

$$(9) \quad \tilde{V}'\tilde{V} = \tilde{I},$$

which imply  $\tilde{\Sigma} = \tilde{V}\tilde{\Lambda}\tilde{V}$  and  $\tilde{I} = \tilde{V}\tilde{V}'$ . We can refer the matrix of independent variables to this coordinate system. Then

$$(10) \quad \tilde{Z} = \tilde{V}\tilde{G},$$

where

$$(11) \quad \tilde{G}' = (\tilde{g}_1, \dots, \tilde{g}_T),$$

and  $\tilde{g}_t$  is a  $p$ -component vector,  $t = 1, \dots, T$ . The two covariance matrices depend on three matrices involving  $\tilde{Z}$  and  $\tilde{\Sigma}$ . These can be written in terms of  $\tilde{\Lambda}$  and  $\tilde{G}$  as

$$(12) \quad \tilde{Z}'\tilde{Z} = \tilde{G}'\tilde{V}'\tilde{V}\tilde{G} = \tilde{G}'\tilde{G} = \sum_{t=1}^T \tilde{g}_t\tilde{g}_t',$$

$$(13) \quad \underset{\sim}{Z}' \underset{\sim}{\Sigma} \underset{\sim}{Z} = \underset{\sim}{G}' \underset{\sim}{V}' \underset{\sim}{\Sigma} \underset{\sim}{V} \underset{\sim}{G} = \underset{\sim}{G}' \underset{\sim}{\Lambda} \underset{\sim}{G} = \sum_{t=1}^T \lambda_t \underset{\sim}{g}_t \underset{\sim}{g}_t' ,$$

$$(14) \quad \underset{\sim}{Z}' \underset{\sim}{\Sigma}^{-1} \underset{\sim}{Z} = \underset{\sim}{G}' \underset{\sim}{V}' \underset{\sim}{\Sigma}^{-1} \underset{\sim}{V} \underset{\sim}{G} = \underset{\sim}{G}' \underset{\sim}{\Lambda}^{-1} \underset{\sim}{G} = \sum_{t=1}^T \frac{1}{\lambda_t} \underset{\sim}{g}_t \underset{\sim}{g}_t' .$$

The columns of  $\underset{\sim}{V}$  are characteristic vectors of  $\underset{\sim}{\Sigma}^{-1}$  corresponding to roots which are the reciprocals of the characteristic roots of  $\underset{\sim}{\Sigma}$ . We shall follow these matrices along.

The characteristic roots may not all be different. Let us indicate the multiplicity of the roots by writing the diagonal matrix  $\underset{\sim}{\Lambda}$  in the partitioned form

$$(15) \quad \underset{\sim}{\Lambda} = \begin{pmatrix} v_1 I & 0 & \dots & 0 \\ 0 & v_2 I & \dots & 0 \\ \vdots & \vdots & & \vdots \\ 0 & 0 & \dots & v_H I \end{pmatrix} ,$$

where  $v_1 > v_2 > \dots > v_H (> 0)$  are the different characteristic roots. The orders of the diagonal blocks are the multiplicities of the corresponding roots, say  $m_1, m_2, \dots, m_H$   $\left( \sum_{h=1}^H m_h = T \right)$ . We partition  $\underset{\sim}{V}$  and  $\underset{\sim}{G}$  similarly,

$$(16) \quad \underset{\sim}{V} = \left( \underset{\sim}{V}^{(1)}, \underset{\sim}{V}^{(2)}, \dots, \underset{\sim}{V}^{(H)} \right) ,$$

$$(17) \quad \underset{\sim}{G} = \begin{pmatrix} \underset{\sim}{G}^{(1)} \\ \underset{\sim}{G}^{(2)} \\ \vdots \\ \underset{\sim}{G}^{(H)} \end{pmatrix} .$$



Now let us go back to the matrices we considered previously, and express them in these new terms.  $\tilde{Z}$  is written as

$$(18) \quad \tilde{Z} = \sum_{h=1}^H \tilde{V}^{(h)} \tilde{G}^{(h)} .$$

The three matrices appearing in the covariance matrices are

$$(19) \quad \tilde{Z}'\tilde{Z} = \sum_{h=1}^H \tilde{G}^{(h)'} \tilde{G}^{(h)} ,$$

$$(20) \quad \tilde{Z}'\tilde{\Sigma}\tilde{Z} = \sum_{h=1}^H v_h \tilde{G}^{(h)'} \tilde{G}^{(h)} ,$$

$$(21) \quad \tilde{Z}'\tilde{\Sigma}^{-1}\tilde{Z} = \sum_{h=1}^H \frac{1}{v_h} \tilde{G}^{(h)'} \tilde{G}^{(h)} .$$

The definition of a submatrix of  $\tilde{V}$  may have some indeterminacy in it. We can replace  $\tilde{V}^{(h)}$  by  $\tilde{V}^{(h)} \tilde{Q}^{(h)}$  and replace  $\tilde{G}^{(h)}$  by  $\tilde{Q}^{(h)'} \tilde{G}^{(h)}$ , where  $\tilde{Q}^{(h)}$  is an orthogonal matrix of order  $m_h$ . Such a transformation leaves each of the last four equations invariant.

Theorem 1 shall be shown to be equivalent to the following theorem:

Theorem 2.  $\mathcal{L}(\tilde{b}) = \mathcal{L}(\tilde{b}^*)$  if and only if

$$(22) \quad \sum_{h=1}^H \rho(\tilde{G}^{(h)}) = p ,$$

where  $\rho(\tilde{G}^{(h)})$  denotes the rank of  $\tilde{G}^{(h)}$ .

In order to simplify the study of the conditions for the equality of the covariance matrices, it is convenient to transform the matrices again. Let  $\tilde{P}$  be a nonsingular matrix such that

$$(23) \quad \tilde{P}'(\tilde{Z}'\tilde{Z})\tilde{P} = \tilde{I} ,$$

$$(24) \quad \tilde{P}'(\tilde{Z}'\tilde{\Sigma}\tilde{Z})\tilde{P} = \tilde{D} ,$$

where  $\tilde{D}$  is a diagonal matrix with  $d_{11} \geq d_{22} \geq \dots \geq d_{pp} \geq 0$ . [These are the characteristic roots of  $\tilde{Z}'\tilde{\Sigma}\tilde{Z}(\tilde{Z}'\tilde{Z})^{-1}$ .] Let us also make the transformation of the other matrix,  $\tilde{P}'(\tilde{Z}'\tilde{\Sigma}^{-1}\tilde{Z})\tilde{P}$ . The covariance matrix of  $\tilde{P}^{-1}\tilde{b}$  is the inverse of this last matrix. The covariance matrix of  $\tilde{P}^{-1}\tilde{b}^*$  is  $\tilde{D}$ . (This can be seen from the original expression for the covariance matrix of  $\tilde{b}^*$ , (6), by multiplication on the left by  $\tilde{P}^{-1}$  and on the right by  $\tilde{P}'^{-1}$  and with use of the properties of the matrices we have just discussed.) The question of equality of the original covariance matrices has now been reduced to the problem of when the covariance matrix of  $\tilde{P}^{-1}\tilde{b}$  is  $\tilde{D}$ .

The three matrices in  $\mathcal{C}(\tilde{b})$  and  $\mathcal{C}(\tilde{b}^*)$  can be written

$$(25) \quad \tilde{I} = \tilde{P}'\tilde{Z}'\tilde{Z}\tilde{P} = \sum_{h=1}^H \tilde{C}^{(h)} ,$$

$$(26) \quad \tilde{D} = \tilde{P}'\tilde{Z}'\tilde{\Sigma}\tilde{Z}\tilde{P} = \sum_{h=1}^H v_h \tilde{C}^{(h)} ,$$

$$(27) \quad \tilde{P}'\tilde{Z}'\tilde{\Sigma}^{-1}\tilde{Z}\tilde{P} = \sum_{h=1}^H \frac{1}{v_h} \tilde{C}^{(h)} ,$$

where  $\tilde{C}^{(h)} = \tilde{P}'\tilde{G}^{(h)'}\tilde{G}^{(h)}\tilde{P}$ . Note that  $\rho(\tilde{C}^{(h)}) = \rho(\tilde{G}^{(h)})$ . Let us consider the diagonal elements of each of the last three equations.

They are

$$(28) \quad 1 = \sum_{h=1}^H c_{ii}^{(h)} ,$$

$$(29) \quad d_{ii} = \sum_{h=1}^H v_h c_{ii}^{(h)},$$

$$(30) \quad \sum_{h=1}^H \frac{1}{v_h} c_{ii}^{(h)}.$$

Since the matrix  $\tilde{C}^{(h)}$  is positive semidefinite, each diagonal element is nonnegative. For each  $i$  the sum of these nonnegative components is 1; hence, the elements in the  $i$ -th diagonal position can be considered as probabilities. Let  $X_i$  be a random variable that takes on the value  $v_h$  with probability  $c_{ii}^{(h)}$ ,  $h = 1, \dots, H$ . Then  $d_{ii}$  is the expected value of this random variable. The last expression is the expected value of the reciprocal of this positive random variable. If the two covariance matrices are to be the same, the  $i$ -th diagonal element of the last matrix must be the reciprocal of that diagonal element of the second matrix. Thus, the random variable just defined can take on only one value with probability 1. (This is basically the condition for equality in the Cauchy-Schwarz inequality.) This implies that for each  $i$ ,  $c_{ii}^{(h)} = 1$  for one index  $h$  and is 0 for other values of  $h$  because the  $v_h$  are distinct. These facts imply that the diagonal elements of the matrices  $\tilde{C}^{(h)}$  are 1's and 0's. The matrices  $\tilde{C}^{(h)}$  have diagonal elements as follows:

$$(31) \quad \tilde{C}^{(1)} = \begin{bmatrix} 1 & & & & & \\ & \ddots & & & & \\ & & 1 & & & \\ & & & 0 & & \\ & & & & \ddots & \\ & & & & & 0 \\ & & & & & & \ddots \\ & & & & & & & 0 \end{bmatrix}, \quad \tilde{C}^{(2)} = \begin{bmatrix} 0 & & & & & \\ & \ddots & & & & \\ & & 0 & & & \\ & & & 1 & & \\ & & & & \ddots & \\ & & & & & 1 \\ & & & & & & \ddots \\ & & & & & & & 0 \end{bmatrix}, \dots$$

If a matrix  $\tilde{C}^{(h)}$  has 1 in the  $i$ -th diagonal position, the other matrices have 0 in that position. (Then  $d_{ii} = v_h$ . Since the  $v_h$ 's and  $d_{ii}$ 's are numbered in descending order, the 1's in  $\tilde{C}^{(1)}$  are in the upper left-hand corner, etc.)

Some matrices may only have 0's on the main diagonal. Since  $\tilde{C}^{(h)}$  is positive semidefinite, a diagonal element of 0 implies that the entire corresponding row and column are 0. Thus

$$(32) \quad \tilde{C}^{(1)} = \begin{bmatrix} 1 & & & 0 \\ & \ddots & & \\ & & 1 & \\ 0 & & & 0 \end{bmatrix}, \quad \tilde{C}^{(2)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & & \ddots & \\ 0 & 0 & 0 & 1 \end{bmatrix}, \dots$$

Since the  $\tilde{C}^{(h)}$ 's sum to  $\tilde{I}$ , and the nonzero blocks are not overlapping

$$(33) \quad \tilde{C}^{(1)} = \begin{bmatrix} \tilde{I} & 0 \\ 0 & 0 \end{bmatrix}, \quad \tilde{C}^{(2)} = \begin{bmatrix} 0 & 0 & 0 \\ 0 & \tilde{I} & 0 \\ 0 & 0 & 0 \end{bmatrix}, \dots$$

We have then  $\tilde{C}^{(1)}$  with an identity in the upper left-hand corner and so on. The rank of each  $\tilde{C}^{(h)}$  is equal to the number of diagonal elements that are 1. Thus, the sum of the ranks is equal to  $p$ . Therefore, the equality of the covariance matrices implies that the sum of the ranks is  $p$ .

The converse can be obtained by use of Cochran's theorem. (See Lemma 7.4.1 of [2], for example.) However, we shall use a simplified proof of a generalization of one part of Cochran's Theorem due to

Styan [8]. We assume the sum of the ranks of the  $\tilde{C}^{(h)}$ 's is  $p$ . Let the nonnull  $\tilde{C}^{(h)}$ 's be  $\tilde{L}_1, \dots, \tilde{L}_K$  ( $K \leq p$ ) and let the ranks of these matrices be  $r_1, \dots, r_K$ , respectively. Then  $\tilde{L}_j$  can be written  $\tilde{A}_j' \tilde{A}_j$ , where  $\tilde{A}_j$  is  $r_j \times p$ ,  $j = 1, \dots, K$ . Let  $\tilde{U}$  be the diagonal matrix with  $j$ -th diagonal blocks of order  $r_j$  consisting of  $u_j \tilde{I}$ , respectively, where  $u_j$  is the  $j$ -th value of  $v_1, \dots, v_H$  corresponding to a nonnull  $\tilde{C}^{(h)}$ ,  $j = 1, \dots, K$ . Let

$$(34) \quad \tilde{A} = \begin{pmatrix} \tilde{A}_1 \\ \vdots \\ \tilde{A}_K \end{pmatrix}.$$

Then (25) and (26) are

$$(35) \quad \tilde{I} = \tilde{A}' \tilde{A},$$

$$(36) \quad \tilde{D} = \tilde{A}' \tilde{U} \tilde{A}.$$

Equation (35) shows that  $\tilde{A}$  is orthogonal as  $\sum_{j=1}^K r_j = p$ , and so it follows from (36) that

$$(37) \quad \tilde{D}^{-1} = \tilde{A}' \tilde{U}^{-1} \tilde{A},$$

which is (27). Since

$$(38) \quad \sum_{h=1}^H \rho(\tilde{C}^{(h)}) = \sum_{j=1}^K r_j = p$$

Theorem 2 is proved. (That equality of covariance matrices implies the rank condition can be proved by the method used in the converse, but it does not generalize directly to the case of stationary residuals.)

As was indicated earlier,  $\tilde{G}^{(h)}$  in  $\tilde{Z} = \sum_{h=1}^H \tilde{V}^{(h)} \tilde{G}^{(h)}$  can be replaced by  $\tilde{Q}^{(h)'} \tilde{G}^{(h)}$  where  $\tilde{Q}^{(h)}$  is orthogonal. In particular,  $\tilde{Q}^{(h)}$  can be chosen so that  $\tilde{G}^{(h)}$  has as many nonzero rows as its rank. (For the nonnull  $\tilde{C}^{(h)}$ 's or  $\tilde{G}^{(h)}$ 's, the resulting matrices are  $\tilde{A}_1, \dots, \tilde{A}_K$ .) This proves Theorem 1 for the finite-dimensional case.

### 3. Large-Sample Theory for Stationary Residuals

We now turn to the problem involving stationary time series. The elements of the covariance matrix of  $\tilde{y}$  are

$$(39) \quad \sigma_{st} = \sigma(s - t) = \int_{-\pi}^{\pi} e^{i(s-t)\lambda} f(\lambda) d\lambda ,$$

where  $f(\lambda)$  is the spectral density, which is assumed to exist. Also we assume that the spectral density satisfies the inequalities

$$(40) \quad 0 < \frac{m}{2\pi} \leq f(\lambda) \leq \frac{M}{2\pi} ,$$

when  $m$  and  $M$  are some positive constants. In developing the asymptotic theory I shall not attempt to state all of the conditions. (They are given in Section 10.2.3 of [3].) We write

$$(41) \quad f(\lambda) = \frac{1}{2\pi} \sum_{h=-\infty}^{\infty} e^{i\lambda h} \sigma(h) .$$

Let the diagonal matrix  $\tilde{D}_T$  be defined by

$$(42) \quad \text{diag} (\tilde{Z}'\tilde{Z}) = \text{diag} \left( \tilde{D}_T^2 \right) ,$$

where we take the positive square roots. Since we are interested in  $T \rightarrow \infty$ , we shall use the index  $T$  when convenient to emphasize that we

have a sequence of estimates. The suitable normalization of the estimates is multiplication by this matrix  $\tilde{D}_T$ . We consider the limits of the covariance matrices of  $\tilde{D}_T b$  and of  $\tilde{D}_T b^*$ . The question is what are necessary and sufficient conditions on the independent variables and the spectral density such that

$$(43) \quad \lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T b) = \lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T b^*) .$$

Let

$$(44) \quad \tilde{Z}' = (\tilde{z}_1, \dots, \tilde{z}_T) .$$

Consider the sum on  $t$  of  $\tilde{z}_{t+h} \tilde{z}_t'$  and multiply on each side by  $\tilde{D}_T^{-1}$  to obtain the matrix of lagged correlations of order  $h$ . Let the limit of this matrix as  $T \rightarrow \infty$  be

$$(45) \quad \tilde{R}(h) = \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \sum_t \tilde{z}_{t+h} \tilde{z}_t' \tilde{D}_T^{-1} .$$

We assume that these limits exist for  $t = 0, \pm 1, \pm 2, \dots$ . Then this sequence of matrices has the spectral representation

$$(46) \quad \tilde{R}(h) = \int_{-\pi}^{\pi} e^{i\lambda h} d\tilde{M}(\lambda) ,$$

where  $\tilde{M}(\lambda)$  has complex-valued elements, is Hermitian, and has increments that are positive semidefinite.

We shall now consider the limits of the covariance matrices of the normalized estimates. Those covariance matrices involve the limits of the matrices  $\tilde{D}_T^{-1} \tilde{Z}' \tilde{Z} \tilde{D}_T^{-1}$ ,  $\tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1}$ , and  $\tilde{D}_T^{-1} \tilde{Z} \tilde{\Sigma}^{-1} \tilde{Z} \tilde{D}_T^{-1}$ . In fact,

$$(47) \quad \lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T \tilde{b}) = \lim_{T \rightarrow \infty} (\tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z} \tilde{D}_T^{-1})^{-1} ,$$

$$(48) \quad \lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T \tilde{b}^*) = \tilde{R}^{-1}(0) \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} \tilde{R}^{-1}(0) .$$

The second matrix is

$$(49) \quad \begin{aligned} \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \sum_{h=-(T-1)}^{T-1} \sum_t \tilde{z}_{t+h} \tilde{z}_t' \sigma(h) \tilde{D}_T^{-1} &= \sum_{h=-\infty}^{\infty} \tilde{R}(h) \sigma(h) \\ &= \int_{-\pi}^{\pi} \sum_{h=-\infty}^{\infty} \sigma(h) e^{i\lambda h} d\tilde{M}(\lambda) \\ &= \int_{-\pi}^{\pi} 2\pi f(\lambda) d\tilde{M}(\lambda) . \end{aligned}$$

Of course, these operations need to be justified to give a rigorous proof, but that requires considerable detail. The full proof is given in section 10.2.3 of my book [3] and is along the lines indicated by Grenander and Rosenblatt [5]. The three matrices we are interested in can be written

$$(50) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{Z} \tilde{D}_T^{-1} = \int_{-\pi}^{\pi} d\tilde{M}(\lambda) ,$$

$$(51) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} = 2\pi \int_{-\pi}^{\pi} f(\lambda) d\tilde{M}(\lambda) ,$$

$$(52) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z} \tilde{D}_T^{-1} = \int_{-\pi}^{\pi} \frac{1}{2\pi f(\lambda)} d\tilde{M}(\lambda) .$$

The derivation for the third matrix is an involved demonstration also given in [3]. These three expressions are the analogues of (12), (13), and (14) in the finite-dimensional case. Carrying the analogy to the finite-dimensional case further, we shall write these integrals in another manner to resemble (19), (20), and (21). Let



$$(53) \quad S(u) = \{\lambda \mid 2\pi f(\lambda) \leq u\}, \quad m \leq u \leq M,$$

$$(54) \quad \tilde{T}(u) = \int_{S(u)} d\tilde{M}(\lambda).$$

The component functions of  $\tilde{T}(u)$  are real. Then our three matrices can be written as

$$(55) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{Z} \tilde{D}_T^{-1} = \int_m^M d\tilde{T}(u),$$

$$(56) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} = \int_m^M u d\tilde{T}(u),$$

$$(57) \quad \lim_{T \rightarrow \infty} \tilde{D}_T^{-1} \tilde{Z} \tilde{\Sigma}^{-1} \tilde{Z}' \tilde{D}_T^{-1} = \int_m^M \frac{1}{u} d\tilde{T}(u).$$

Similar to the finite-dimensional case we let  $\tilde{P}$  be a nonsingular matrix such that

$$(58) \quad \tilde{P}' \int d\tilde{T}(u) \tilde{P} = \tilde{I},$$

$$(59) \quad \tilde{P}' \int u d\tilde{T}(u) \tilde{P} = \tilde{D},$$

where  $\tilde{D}$  is diagonal and  $d_{11} \geq d_{22} \geq \dots \geq d_{pp} \geq 0$ . The same transformation is applied to the third matrix,  $\int_m^M u^{-1} d\tilde{M}(u)$ , which is the inverse of  $\lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T b)$ . The other limiting covariance matrix is  $\lim_{T \rightarrow \infty} \mathcal{C}(\tilde{D}_T b^*) = \tilde{D}$ .

If we let

$$(60) \quad \tilde{L}(u) = \tilde{P}' \tilde{T}(u) \tilde{P},$$

then the three matrices of interest are

$$(61) \quad \tilde{I} = \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \int_m^M d\tilde{L}(u) ,$$

$$(62) \quad \tilde{D} = \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \int_m^M u d\tilde{L}(u) ,$$

$$(63) \quad \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \int_m^M \frac{1}{u} d\tilde{L}(u) .$$

A diagonal element of  $\tilde{L}(u)$ , say  $\ell_{ii}(u)$ , has the properties of a cumulative distribution function. The corresponding diagonal elements of (62) and (63) are  $\int_m^M u d\ell_{ii}(u)$  and  $\int_m^M u^{-1} d\ell_{ii}(u)$ , which are the expected values of the random variable with this distribution and its reciprocal. Thus, if the two limiting covariance matrices are equal, the matrix (63) is the inverse of (62) and

$$(64) \quad \int_m^M u d\ell_{ii}(u) = \left[ \int_m^M \frac{1}{u} d\ell_{ii}(u) \right]^{-1} ;$$

this implies that  $\ell_{ii}(u)$  has one point of increase and the increase is 1 at this point. Let the points of increase be  $u_1 > u_2 \dots > u_K > 0$ , and let  $L_j$  be the increase of  $\tilde{L}(u)$  at  $u_j$ ,  $j = 1, \dots, K$ . Then the three matrices can be written

$$(65) \quad \tilde{I} = \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \sum_{j=1}^K \tilde{L}_j ,$$

$$(66) \quad \tilde{D} = \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \sum_{j=1}^K u_j \tilde{L}_j ,$$

$$(67) \quad \lim_{T \rightarrow \infty} \tilde{P}' \tilde{D}_T^{-1} \tilde{Z}' \tilde{\Sigma}^{-1} \tilde{Z} \tilde{D}_T^{-1} \tilde{P} = \sum_{j=1}^K \frac{1}{u_j} \tilde{L}_j .$$

We are now back to the same forms that we had for the finite-dimensional case, (25), (26), (27). The only difference is that in the earlier

case we had not culled out the vacuous matrices  $\tilde{C}^{(h)}$ . From this point on the reasoning is the same. The matrices  $\tilde{L}_1, \tilde{L}_2, \dots, \tilde{L}_K$  have the form of (33); that is, the diagonal blocks are  $\tilde{I}$ 's and  $\tilde{O}$ 's and off-diagonal blocks are  $\tilde{O}$ 's.

The converse is similar to the finite-dimensional case. If  $\tilde{L}(u)$  has  $K$  points of increase and the sum of the ranks of the increases is  $p$  (and the increases are positive semidefinite with sum of  $\tilde{I}$  and weighted sum of  $\tilde{D}$ ), then by the previous reasoning, they are of the form (33) and (67) is  $\tilde{D}^{-1}$ . We put these properties in terms of  $\tilde{M}(\lambda)$  and summarize them in a theorem.

Theorem 3. The limiting covariances of  $\tilde{D}_T \tilde{b}$  and  $\tilde{D}_T \tilde{b}^*$  are identical if and only if  $f(\lambda)$  takes on no more than  $p$  values on the set of  $\lambda$  for which  $\tilde{M}(\lambda)$  increases and the sum of the ranks of  $\int \tilde{M}(\lambda)$  over the sets of  $\lambda$  for which  $f(\lambda)$  takes on these values is  $p$ .

The set of  $\lambda$  for which  $\tilde{M}(\lambda)$  increases is called the spectrum of  $\tilde{M}(\lambda)$ . The sets of  $\lambda$  for which  $f(\lambda)$  assumes its values are called the elements of the spectrum. The properties of  $\tilde{L}_1, \dots, \tilde{L}_K$  (idempotent and orthogonal) determine these sets; Grenander and Rosenblatt used them, though indirectly.

When the residuals are uncorrelated,  $f(\lambda) = \sigma(0)/(2\pi)$  and the conditions of Theorem 3 are satisfied. However, we may be interested in conditions on the independent variables also which insure that least squares be asymptotically efficient regardless of  $f(\lambda)$ .

Theorem 4. The limiting covariances of  $D_{\sim T} b$  and  $D_{\sim T} b^*$  are identical for all stationary processes with spectral densities which are bounded and bounded away from 0 if and only if  $\tilde{M}(\lambda)$  increases at not more than  $p$  values of  $\lambda$ ,  $0 \leq \lambda \leq \pi$ , and the sum of the ranks of the increase in  $\tilde{M}(\lambda)$  is  $p$ .

If the number of points at which  $\tilde{M}(\lambda)$  increases is at most  $p$  ( $0 \leq \lambda \leq \pi$ ), then the spectrum of  $\tilde{M}(\lambda)$  consists of these  $p$  points and their corresponding negative values. The spectral density (which is symmetric) can then take on at most  $p$  values, namely, its values at these  $p$  points ( $0 \leq \lambda \leq \pi$ ). On the other hand if  $\tilde{M}(\lambda)$  increases at more than  $p$  points ( $0 \leq \lambda \leq \pi$ ) then an  $f(\lambda)$  can be constructed so that it takes on more than  $p$  points.

An example of independent variables  $\{z_{jt}\}$  such that  $\tilde{M}(\lambda)$  has one point of increase is  $z_{jt} = t^{j-1}$ ,  $j = 1, \dots, p$ ,  $t = 1, 2, \dots$ ; the jump is at 0 and the increase in  $\tilde{M}(\lambda)$  at  $\lambda = 0$  is a positive definite matrix

$$(68) \quad \tilde{M}_0 = \begin{pmatrix} \sqrt{2j-1} & \sqrt{2k-1} \\ j+k-1 \end{pmatrix}.$$

In this case  $\tilde{R}(h) = \tilde{M}_0$ ,  $h = 0, \pm 1, \dots$ . If

$$(69) \quad z_t = \alpha_0 + \sum_{j=1}^H (\alpha_j \cos v_j t + \beta_j \sin v_j t),$$

then  $\tilde{M}(\lambda)$  has an increase of rank 1 at  $\lambda = 0$  and an increase of rank 2 as  $\lambda = v_j$  (with  $0 < v_j < \pi$ ),  $j = 1, \dots, H$ . In these examples the spectral distribution function of each independent variable

is a pure jump function, which can be considered as the opposite of a density. Trigonometric functions act like characteristic vectors of a covariance matrix in the sense that they are involved in spectral representation. Comparison of  $\tilde{\Sigma} = \tilde{V}\tilde{\Lambda}\tilde{V}'$  and (39) suggests that columns of  $\tilde{V}$  correspond to functions  $e^{i\lambda s}$ , the diagonal components of  $\tilde{\Lambda}$  correspond to the values of  $2\pi f(\lambda)$ , and summation with respect to the index of diagonal components of  $\tilde{\Lambda}$  corresponds to integration with respect to  $\lambda/(2\pi)$ . The analogue of  $\tilde{V}'\tilde{\Sigma}\tilde{V} = \tilde{\Lambda}$  is (41), which involves a limiting procedure.

## REFERENCES

- [ 1] T. W. ANDERSON, "On the theory of testing serial correlation," Skandinavisk Aktuarietidskrift, Vol. 31 (1948), pp. 88-116.
- [ 2] T. W. ANDERSON, An Introduction to Multivariate Statistical Analysis, New York, Wiley, 1958.
- [ 3] T. W. ANDERSON, The Statistical Analysis of Time Series, New York, Wiley, 1971.
- [ 4] ULF GRENANDER, "On the estimation of regression coefficients in the case of an autocorrelated disturbance," Annals of Mathematical Statistics, Vol. 25 (1954), pp. 252-272.
- [ 5] ULF GRENANDER and MURRAY ROSENBLATT, Statistical Analysis of Stationary Time Series, New York, Wiley, 1957.
- [ 6] T. A. MAGNESS and J. B. McGUIRE, "Comparison of least squares and minimum variance estimates of regression parameters," Annals of Mathematical Statistics, Vol. 33 (1962), pp. 462-470.
- [ 7] MURRAY ROSENBLATT, "Some regression problems in time series analysis," Proceedings of the Third Berkeley Symposium on Mathematical Statistics and Probability, Vol. 1, Berkeley and Los Angeles, University of California Press, 1956.
- [ 8] GEORGE P. H. STYAN, "Notes on the distribution of quadratic forms in singular normal variables," Biometrika, Vol. 57 (1970),
- [ 9] G. S. WATSON, "Serial correlation in regression," Mimeo Ser. No. 49, Institute of Statistics, University of North Carolina.
- [10] G. S. WATSON, "Serial correlation in regression analysis. I," Biometrika, Vol. 42 (1955), pp. 326-341.
- [11] G. S. WATSON, "Linear least squares regression," Annals of Mathematical Statistics, Vol. 38 (1967), pp. 1679-1699.
- [12] G. S. WATSON and E. J. HANNAN, "Serial correlation in regression analysis. II," Biometrika, Vol. 43 (1956), pp. 436-448.
- [ 13] GEORGE ZYSKIND, "On canonical forms, non-negative covariance matrices and best and simple least squares linear estimation in linear models," Annals of Mathematical Statistics, Vol. 38 (1967), pp. 1092-1109.

UNCLASSIFIED

## Security Classification

## DOCUMENT CONTROL DATA - R&amp;D

(Security classification of title, body of abstract and indexing annotation must be entered when the overall report is classified)

1. ORIGINATING ACTIVITY (Corporate author) DEPARTMENT OF STATISTICS STANFORD UNIVERSITY STANFORD, CALIF.		2a. REPORT SECURITY CLASSIFICATION	
		2b. GROUP	
3. REPORT TITLE  EFFICIENT ESTIMATION OF REGRESSION COEFFICIENTS IN TIME SERIES			
4. DESCRIPTIVE NOTES (Type of report and inclusive dates) TECHNICAL REPORT			
5. AUTHOR(S) (Last name, first name, initial)  ANDERSON, T. W.			
6. REPORT DATE OCTOBER 1, 1970	7a. TOTAL NO. OF PAGES 20	7b. NO. OF REFS 13	
8a. CONTRACT OR GRANT NO. N00014-67-A-0112-0030	9a. ORIGINATOR'S REPORT NUMBER(S)  Technical Report No. 2		
b. PROJECT NO. NR-042-034	9b. OTHER REPORT NO(S) (Any other numbers that may be assigned this report)		
c.			
d.			
10. AVAILABILITY/LIMITATION NOTICES Unlimited. Reproduction in whole or in part is permitted for any purpose of the United States Government.			
11. SUPPLEMENTARY NOTES		12. SPONSORING MILITARY ACTIVITY OFFICE OF NAVAL RESEARCH Washington, D.C.	
13. ABSTRACT  Problems of efficient estimation are considered in the model in which the T-component observable random vector $y$ has expected value $Z\beta$ , where $Z$ is a $T \times p$ matrix of known numbers of rank $p$ ( $p \leq T$ ) and $\beta$ is a $p$ -component vector of regression coefficients, and (nonsingular) covariance matrix $\Sigma$ . The least squares estimate of $\beta$ is identical to the Markov or Best Linear Unbiased Estimate if and only if the $p$ columns of $Z$ are linearly independent linear combinations of $p$ linearly independent characteristic vectors of $\Sigma$ . The proof uses the covariance matrices of the estimates. When the components constitute time series such that $\Sigma$ corresponds to a stochastic process stationary in the wide sense, the limits of the appropriately normalized covariance matrices of the two estimates are considered as $T \rightarrow \infty$ . They are functions of the matrix-valued spectral distribution function $M(\lambda)$ of the limiting serial correlation sequence of the rows of $Z$ . On the basis of a similar approach it is shown that these two limiting covariance matrices are identical if and only if the spectral density of the process whose covariances are elements of $\Sigma$ takes on at most $p$ values on the set of $\lambda$ for which $M(\lambda)$ increases and the sum of the ranks of $\int dM(\lambda)$ over the sets of $\lambda$ for which the density takes on the value is $p$ . The result holds for all such spectral densities if and only if $M(\lambda)$ increases at not more than $p$ values of $\lambda$ , $0 \leq \lambda \leq \pi$ , and the sum of the ranks of the increases is $p$ .			

DD FORM 1473  
1 JAN 64

UNCLASSIFIED

Security Classification

14. KEY WORDS	LINK A		LINK B		LINK C	
	ROLE	WT	ROLE	WT	ROLE	WT
1. Efficient estimation						
2. Least squares estimates						
3. Best linear unbiased estimates						
4. Time series						
5. Spectral distribution						
6. Asymptotic efficiency						

#### INSTRUCTIONS

**1. ORIGINATING ACTIVITY:** Enter the name and address of the contractor, subcontractor, grantee, Department of Defense activity or other organization (corporate author) issuing the report.

**2a. REPORT SECURITY CLASSIFICATION:** Enter the overall security classification of the report. Indicate whether "Restricted Data" is included. Marking is to be in accordance with appropriate security regulations.

**2b. GROUP:** Automatic downgrading is specified in DoD Directive 5200.10 and Armed Forces Industrial Manual. Enter the group number. Also, when applicable, show that optional markings have been used for Group 3 and Group 4 as authorized.

**3. REPORT TITLE:** Enter the complete report title in all capital letters. Titles in all cases should be unclassified. If a meaningful title cannot be selected without classification, show title classification in all capitals in parentheses immediately following the title.

**4. DESCRIPTIVE NOTES:** If appropriate, enter the type of report, e.g., interim, progress, summary, annual, or final. Give the inclusive dates when a specific reporting period is covered.

**5. AUTHOR(S):** Enter the name(s) of author(s) as shown on or in the report. Enter last name, first name, middle initial. If military, show rank and branch of service. The name of the principal author is an absolute minimum requirement.

**6. REPORT DATE:** Enter the date of the report as day, month, year, or month, year. If more than one date appears on the report, use date of publication.

**7a. TOTAL NUMBER OF PAGES:** The total page count should follow normal pagination procedures, i.e., enter the number of pages containing information.

**7b. NUMBER OF REFERENCES:** Enter the total number of references cited in the report.

**8a. CONTRACT OR GRANT NUMBER:** If appropriate, enter the applicable number of the contract or grant under which the report was written.

**8b, 8c, & 8d. PROJECT NUMBER:** Enter the appropriate military department identification, such as project number, subproject number, system numbers, task number, etc.

**9a. ORIGINATOR'S REPORT NUMBER(S):** Enter the official report number by which the document will be identified and controlled by the originating activity. This number must be unique to this report.

**9b. OTHER REPORT NUMBER(S):** If the report has been assigned any other report numbers (either by the originator or by the sponsor), also enter this number(s).

**10. AVAILABILITY/LIMITATION NOTICES:** Enter any limitations on further dissemination of the report, other than those

imposed by security classification, using standard statements such as:

- (1) "Qualified requesters may obtain copies of this report from DDC."
- (2) "Foreign announcement and dissemination of this report by DDC is not authorized."
- (3) "U. S. Government agencies may obtain copies of this report directly from DDC. Other qualified DDC users shall request through \_\_\_\_\_."
- (4) "U. S. military agencies may obtain copies of this report directly from DDC. Other qualified users shall request through \_\_\_\_\_."
- (5) "All distribution of this report is controlled. Qualified DDC users shall request through \_\_\_\_\_."

If the report has been furnished to the Office of Technical Services, Department of Commerce, for sale to the public, indicate this fact and enter the price, if known.

**11. SUPPLEMENTARY NOTES:** Use for additional explanatory notes.

**12. SPONSORING MILITARY ACTIVITY:** Enter the name of the departmental project office or laboratory sponsoring (paying for) the research and development. Include address.

**13. ABSTRACT:** Enter an abstract giving a brief and factual summary of the document indicative of the report, even though it may also appear elsewhere in the body of the technical report. If additional space is required, a continuation sheet shall be attached.

It is highly desirable that the abstract of classified reports be unclassified. Each paragraph of the abstract shall end with an indication of the military security classification of the information in the paragraph, represented as (TS), (S), (C), or (U).

There is no limitation on the length of the abstract. However, the suggested length is from 150 to 225 words.

**14. KEY WORDS:** Key words are technically meaningful terms or short phrases that characterize a report and may be used as index entries for cataloging the report. Key words must be selected so that no security classification is required. Identifiers, such as equipment model designation, trade name, military project code name, geographic location, may be used as key words but will be followed by an indication of technical context. The assignment of links, roles, and weights is optional.